



STRATUS CONSULTING

**Greenhouse Gas Emissions
from Fossil Energy Extracted
from Federal Lands and Waters:
An Update
Final**

Prepared for:

The Wilderness Society
1615 M Street NW
Washington, DC 20036

**Greenhouse Gas Emissions
from Fossil Energy Extracted
from Federal Lands and Waters:
An Update
Final**

Prepared for:

The Wilderness Society
1615 M Street NW
Washington, DC 20036

Prepared by:

Stratus Consulting Inc.
PO Box 4059
Boulder, CO 80306-4059
303-381-8000

1920 L St. NW, Suite 420
Washington, DC 20036

Contacts:

Heidi Ries
Carolyn Wagner

September 17, 2014

Confidential

SC13598

1. Introduction

In April 2011, the Council on Environmental Quality (CEQ) released the *Greenhouse Gas Emissions Inventory for the Federal Government: 2010 Data* (CEQ, 2011). This report presented the total estimated greenhouse gas (GHG) emissions associated with federal government agency operations, including emissions from building electricity and water consumption, employee travel, and numerous other activities. However, the inventory did not account for emissions associated with a range of activities subject to federal agency oversight but conducted by private entities. Such activities include exploration, production, and development of fossil fuel resources on federal lands¹ by private sector leaseholders. Accounting for these activities could increase the federal government's total emissions of GHGs substantially.

In 2012, Stratus Consulting produced a report on behalf of The Wilderness Society entitled *Greenhouse Gas Emissions from Fossil Energy Extracted from Federal Lands and Waters* in which we developed a preliminary estimate of the magnitude of ultimate GHG emissions² associated with fossil fuels for the years 2008–2010 – for example, onshore and offshore oil and natural gas, natural gas liquids, coal, and coalbed methane – that leaseholders³ extract from federal lands. The report also included an overview of many types of indirect emissions that could result from fossil fuel extraction, such as emissions from transporting fossil fuels from federal lands to refineries. The report did not address GHG emissions resulting from wind, solar, geothermal, or biomass activities.⁴

1. The term “federal lands” includes all offshore areas that the federal government leases to private companies, as well as onshore lands in this report.

2. Ultimate GHG emissions refer to carbon dioxide (CO₂), methane, and nitrous oxide emissions that result from fossil fuel extracted from federal lands (i.e., the quantity of gas emitted if an amount of a fossil fuel was combusted downstream in various applications, such as coal-fired power plants).

3. We focus on GHG emissions resulting from the extraction of fossil fuels from federal lands by private companies holding leases for these purposes, even in instances where the federal government holds subsurface rights but does not hold surface rights. As such, this report does not account for emissions resulting from the extraction of fossil fuels in instances where the federal government owns the surface rights but private or other government entities own the subsurface rights.

4. We cover a subset of energy extraction activities on federal lands and waters. We focus on fossil fuel extraction from federal lands, where the U.S. Department of the Interior (DOI) manages subsurface leasing. The DOI manages subsurface leasing for all federal lands, including public lands under the jurisdiction of the Bureau of Land Management (BLM) and other lands under the jurisdiction of other federal surface management agencies. The data that we collected from DOI sources for this analysis include information on all leases that the DOI manages.

Our current report provides an updated analysis of the magnitude of GHG emissions currently accounted for from federal lands and waters. Specifically, we add emissions estimates for calendar year 2012 using the same methodology utilized in the 2012 analysis.⁵ We also include an appendix characterizing indirect emissions from natural gas and oil production. We focus on methane emissions from natural gas and oil production to highlight the importance of these emissions and their contribution to climate change. These emissions also highlight the growing importance of unconventional – fracking and tight gas – natural gas production and venting and flaring during oil production. Methane is 25 times more potent than CO₂ as a GHG, and emissions from natural gas and oil production could be significant.

1.1 Background

Federal lands and waters supply a considerable portion of the nation’s fossil energy resources, including approximately 42% of produced coal, 26% of crude oil, and 13% of natural gas.⁶ Rather than developing, extracting, transporting, and converting these resources directly, federal agencies typically operate through permits, leases, contracts, and concessions to private sector firms.

Executive Order (E.O.) 13514, *Federal Leadership in Environmental, Energy, and Economic Performance* (74 Federal Register 52117, October 9, 2009; The White House Office of the Press Secretary, 2009), directed federal agencies to meet a range of energy, water, and waste-reduction targets. The E.O. included requirements that federal agencies inventory, report, and adopt targets for reducing their direct and indirect GHG emissions. The CEQ developed the *Federal Greenhouse Gas Accounting and Reporting Guidance* (CEQ, 2010b) to provide federal agencies with clear information on which emissions fall within the scope of E.O. 13514 and how to calculate these emissions. The report included guidance on how to quantify emissions resulting from a range of source categories. Table 1 lists the GHG emissions source categories included in the guidance and ultimately accounted for in the federal government inventory. Federal government agencies reported their emissions for 2010 and made them public in April 2011.

Total U.S. GHG emissions and emissions related to energy consumption

The U.S. Environmental Protection Agency (EPA) produces an annual GHG inventory every year that provides estimates of the nation’s source- and sector-specific emissions of CO₂, methane (CH₄), nitrous oxide (N₂O), and other GHGs. According to EPA’s most recent (2014) inventory, GHG emissions in the United States in 2012, the most recent year for which data were available, totaled approximately 6,526 million metric tons of carbon dioxide equivalent (MMTCO₂e). Energy-related GHG emissions totaled 5,499 MMTCO₂e.

Source: U.S. EPA, 2014b.

5. Our current report draws heavily on the 2012 report for structure, methodology, and presentation.

6. We collected data for fossil fuel produced from federal lands for this analysis (BOEMRE, 2014; ONRR, 2014b), and data on total fossil fuel production for the entire United States (EIA, 2014b, 2014c, 2014d).

Table 1. Sources of emissions accounted for in the *Greenhouse Gas Emissions Inventory for the Federal Government: 2010 Data*, released by the CEQ in April 2011

On-site fuel combustion
Non-highway vehicles, aircraft, ships, and equipment
Passenger fleet vehicles
Fluorinated gases and other fugitive emissions
On-site wastewater treatment
On-site landfills, solid waste facilities, and incinerators
Manufacturing and industrial process emissions
Purchased electricity
Purchased biomass energy
Purchased steam and hot water
Purchased chilled water
Purchased combined heat and power, electricity, steam, and hot water
Transmission and distribution losses from purchased electricity
Federal employee business air travel
Federal employee business ground travel
Federal employee commuting
Off-site wastewater treatment
Off-site solid waste disposal

Source: CEQ, 2011.

In the report *Greenhouse Gas Emissions Inventory for the Federal Government: 2010 Data*, the CEQ estimated that GHG emissions from federal government operations totaled approximately 66.4 MMTCO₂e in 2010. According to the inventory, DOI, which is responsible for managing a significant majority of federal fossil fuel leases, contributed approximately 1.3 MMTCO₂e (CEQ, 2011). The CEQ guidance did not include requirements for reporting emissions associated with activities, including fossil fuel extraction on federal lands, which are subject to federal agency oversight but that private entities conduct. Federal government agencies, therefore, did not estimate these emissions in the reports they compiled for the inventory.⁷

7. Federal agencies are required to account for GHG emissions under other obligations. For example, federal agencies are obligated under the National Environmental Policy Act (NEPA), 42 U.S.C. § 4321, et seq., to analyze the impacts of major federal actions that have significant impacts on the environment, including the impacts on global climate change. However, emissions resulting from federal leasing are not explicitly addressed in the NEPA guidance for federal agencies (CEQ, 2010a). Agencies within the DOI are additionally bound by Secretarial Order 3289, which requires the analysis of potential climate change effects when undertaking planning, setting priorities for scientific research, developing management plans, and making major decisions regarding the potential use of resources.

Our first report on behalf of The Wilderness Society, entitled *Greenhouse Gas Emissions from Fossil Energy Extracted from Federal Lands and Waters*, estimated that ultimate downstream GHG emissions from federal lands in 2010 could have been more than 20 times the estimate of total GHG emissions in the CEQ report.

1.2 Overview

The remainder of this report is structured as follows:

- ▶ **Section 2 – Methodology.** In this section, we describe the methodology used in the analysis. We include an overview of the process we used to quantify ultimate GHG emissions from fossil fuel extraction on federal lands by private leaseholders.
- ▶ **Section 3 – Results.** Here, we summarize the results of the quantitative analysis of ultimate GHG emissions resulting from the extraction of fossil fuels on federal lands by private leaseholders in 2012. We provide fuel-, location-, and GHG-specific estimates.
- ▶ **Section 4 – Conclusions.** This section presents key points based on the analysis we describe in this report.
- ▶ **References.** This section includes references that we cite in the report.
- ▶ **Appendix – Indirect GHG Emissions from Natural Gas and Oil Production.** Here, we summarize the results of a literature review of indirect GHG emissions from natural gas production. We focus on methane emissions from natural gas and oil production, and quantify emissions for various scenarios involving venting, flaring, and unintentional leaks of natural gas.

2. Methodology

This section describes the methodology used for this analysis. We include an overview of the process used to quantify ultimate GHG emissions from fossil fuel extraction from federal lands by private leaseholders. We discuss data collection including the quantities of fossil fuels extracted from federal lands by private leaseholders, energy flows, and emissions factors. Then we present the steps involved in data processing: converting units, matching quantities of fossil fuels to sectors and emissions factors, and performing calculations.

2.1 Methodology for Quantifying Ultimate GHG Emissions from Fossil Fuel Extraction from Federal Lands by Private Leaseholders

Quantifying ultimate GHG emissions from fossil fuel extraction on federal lands by private leaseholders involved two broad steps: data collection and data processing. This section describes the steps involved in each step.

2.1.1 Data collection

This section describes the data collection portion of the analysis, including the types of data we collected and our data sources.

Quantities of fossil fuels extracted from federal lands by private leaseholders

The primary data collection effort involved gathering information on the quantities of fossil fuels extracted from federal lands by private leaseholders, based on reports by federal government agencies. The data included the quantity of fuel extracted by relevant metric (e.g., barrels of oil, cubic feet of natural gas, and short tons of coal), the year,⁸ and the state in which the lease was held (or region, in the case of offshore data). We collected data for the following fuel categories: onshore and offshore oil, onshore and offshore natural gas, natural gas liquids, coal, and coalbed methane.⁹

We collected data from several DOI sources. Table 2 provides a list of these sources by fuel type.

8. Our analysis used the calendar year as our unit of temporal measure.

9. For onshore oil, onshore natural gas, natural gas liquids, coal, and coalbed methane, quantities were only available in terms of sales volume (as indicated in royalty reports), rather than production volume. In theory, sales volumes for a given year can be smaller or larger than production volumes for that year because companies can vary sales depending on market conditions. For offshore oil and natural gas, quantities were available in terms of production volume.

Table 2. Sources of data on quantities of fossil fuels produced from federal leases

Fossil fuel	Federal government agency and program
Coal and coalbed methane	DOI, BLM – Public Land Statistics: Federal Coal Leases (BLM, 2011) DOI, Office of Natural Resources Revenue (ONRR) – Reported Royalty Revenues (ONRR, 2014b)
Oil, natural gas (onshore), and natural gas liquids	DOI, ONRR – Reported Royalty Revenues (ONRR, 2011, 2014b)
Oil, natural gas (onshore and offshore)	DOI, ONRR – Oil and Natural Gas Report (ONRR, 2014a)
Oil and natural gas (offshore)	DOI, Bureau of Ocean Energy Management, Regulation, and Enforcement – Outer Continental Shelf Oil and Gas Production (BOEMRE, 2014)

Energy flows

We also collected information that enabled us to match the quantities of fuel to specific downstream combustion methods, each of which has a distinct set of emissions factors for key GHGs.¹⁰ We discuss the importance of this fuel type-combustion method matching in the following section. Estimates for 2012 were derived from the Energy Information Administration's (EIA's) *Annual Energy Outlook 2012 with Projections to 2035*, which provided information for the base year, 2012, as well as future projections (EIA, 2009b, 2010, 2011a, 2012a, 2013).

Emissions factors

We identified emissions conversions factors to determine the quantity of GHGs emitted for each unit of fossil fuel extracted. The most widely accepted emissions factors used to calculate GHG emissions were provided in the Intergovernmental Panel on Climate Change's (IPCC's) *IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC, 2006). These guidelines included tables that presented estimates of the kilograms of CO₂, CH₄, and N₂O emitted for every terajoule, or measure of energy content, of energy consumed. These tables included specific fossil fuel- and combustion-type emissions factors. For example, IPCC guidelines provided emissions factors to estimate the kilograms of CH₄, CO₂, and N₂O emitted for every terajoule of energy consumed from burning sub-bituminous coal in a coal-fired power plant for electricity generation.

10. For example, coal combustion for electricity generation has a different set of emissions factors from oil combustion for residential heating.

For the transportation sector, the IPCC presented a number of subsector-specific sets of emissions factors based on the type of fuel used and the combustion process used in each subsector (e.g., for on-road vehicles, IPCC presented different sets of emissions factors for light-duty vehicles and heavy-duty vehicles). We also collected data on the percentages of transportation-related energy that the different subsectors used from EIA's *Annual Energy Outlook 2014 with Projections to 2040* (EIA, 2014a).

For coal, GHG emissions vary depending on the type of coal combusted (e.g., lignite versus sub-bituminous). We collected information on the type of coal extracted in states where private leaseholders extract coal from federal lands from the DOI's *2010 Public Lands Statistics: Federal Coal Leases* (BLM, 2011), which includes reports of the states from which private leaseholders extract coal, and EIA's *Annual Coal Report – 2009* (EIA, 2009a), which provides information on the typical type of coal extracted in each state. Table 3 combines these two sets of information.

Table 3. Coal type by state (states with private leasing on federal lands for coal extraction)

States with private leasing for coal	Dominant coal type
Alabama	Bituminous
Arizona	Bituminous
Colorado	Bituminous
Kentucky	Bituminous
Montana	Sub-bituminous
New Mexico	Sub-bituminous
North Dakota	Lignite
Oklahoma	Bituminous
Ohio	Bituminous
Utah	Bituminous
Washington	Bituminous
Wyoming	Sub-bituminous

Sources: EIA, 2009a; BLM, 2011.

2.1.2 Data processing

To calculate the ultimate GHG emissions from fossil fuel extraction on federal lands by private leaseholders, the data described above were processed in several ways. This section describes the steps we took.

Converting units

The calculation of ultimate GHG emissions involves a number of conversions. For example, the quantities of fossil fuels extracted by private leaseholders from federal lands were reported in volume-based metrics (e.g., short tons, barrels, cubic feet), but the emissions factors provided by the IPCC were characterized in terms of the amount of energy produced by burning the fossil fuel. For this reason, the quantities of fossil fuel extracted were converted into energy-based measures using common conversion factors, such as those available at <http://www.natgas.info/html/natgasunitsconversion.html> (Chandra, 2011). EIA and the International Energy Agency also provided conversion information that was used in this analysis [http://www.eia.gov/kids/energy.cfm?page=about_energy_conversion_calculator-basics (EIA, 2011b) and <http://www.i.e.a.org/stats/unit.asp> (IEA, 2011), respectively].

Matching quantities of fossil fuels to sectors and emissions factors

As noted above, GHG emissions from each fuel type can be determined by multiplying a given quantity of combusted fuel by an emissions factor that is specific to a particular type of combustion. For each of the five sectors identified by the EIA, corresponding sets of combustion-type emissions factors were identified in the *IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC, 2006). For example, the guidelines offered default sets of emissions factors for combustion in stationary power plants, which we deemed appropriate for coal; or combustion in residential settings, which were viewed as appropriate for natural gas.

Information from the EIA was used to classify combustion type by fuel type. These classification “splits” were determined for 2012 by consulting the *Annual Energy Review* publications (EIA, 2012b).

Performing calculations

Equation 1 provides a general illustration of how the above information was combined to produce an estimate of the total amount of GHGs emitted from the combustion of a particular quantity of fossil fuel:

Equation 1. Simplified illustration of equation used to calculate GHG emissions.

$$\begin{aligned}
 & \text{Quantity of GHG emitted (in kilograms)} \\
 &= \sum \text{Quantity of fossil fuel (in terajoules)} \\
 & \quad \times \text{Emissions factor (in kilograms per terajoule)} \times \text{Sector weight}
 \end{aligned}$$

where the sector weight represents the percentage of the quantity of fossil fuel that was consumed in the specific sector (e.g., transportation), thus using the sector-specific set of emissions factors, and the total amount of GHG emitted for a given quantity of fossil fuel is a sum of the calculations for each sector.

Calculations were run to determine the ultimate GHG emissions from federal leases, based on the following:

- ▶ Year
- ▶ GHG (i.e., CO₂, CH₄, or N₂O)
- ▶ Carbon dioxide equivalent (CO₂e; accounting for all three GHGs)¹¹
- ▶ Fossil fuel type
- ▶ Location (onshore versus offshore).

We present the results of these calculations in Section 3.1.

3. Results

This section presents the results of the analysis of ultimate GHG emissions from extraction of fossil fuels from federal lands by private leaseholders. We provide estimates of ultimate GHG emissions by fuel, location, and GHG for 2012.

3.1 Quantification of Ultimate GHG Emissions from Fossil Fuel Extraction from Federal Lands by Private Leaseholders

Table 4 presents estimates of the GHG emissions associated with fossil fuels extracted from federal lands by private leaseholders for 2012.¹² According to this analysis, in 2012 the GHG emissions resulting from the extraction of fossil fuels from federal lands by private leaseholders totaled approximately 1,344 MMTCO₂e.

Figure 1 shows the percentage breakdown of emissions by fossil fuel in 2012.

GHG equivalencies

The GHG emissions from federal lands total 1,344 MMTCO₂e in 2012. These GHG emissions are equivalent to the annual GHG emissions emitted by 283,000,000 passenger vehicles.


Source: U.S. EPA, 2014b.

11. CO₂e were based on the 100-year global warming potentials (GWPs) for these gases, as cited in the IPCC's Fourth Assessment Report on Climate Change (IPCC, 2007).

12. Note that the estimates we present in Tables 5 through 9 are rounded.

Table 4. GHG emissions associated with fossil fuels extracted from federal lands by private leaseholders in 2012

Fossil fuel	Metric tons of CH ₄	Metric tons of CO ₂	Metric tons of N ₂ O	Total MTCO ₂ e
Oil (onshore)	2,999	56,346,510	2,985	57,311,142
Oil (offshore)	11,368	213,556,048	11,315	217,212,051
Natural gas (onshore)	12,358	144,135,798	480	144,587,927
Natural gas (offshore)	8,390	97,851,370	326	98,158,313
Natural gas liquids	11,161	26,134,702	503	26,563,487
Coal	13,080	765,241,950	12,037	769,155,909
Coalbed methane	2,656	30,973,401	103	31,070,559
Total	62,012	1,334,239,779	27,749	1,344,059,388

**Figure 1. Percentage of total MTCO₂e emissions associated with fossil fuels extracted from federal lands by private leaseholders in 2012.**

The estimates above suggest how fossil fuel extraction from federal lands by private leaseholders may contribute to the total amount of GHG emissions included in EPA's national GHG inventory. As noted above, EPA produces a GHG inventory every year that includes estimates of the nation's source- and sector-specific emissions of CO₂, CH₄, N₂O, and other GHGs. According to the 2014 inventory, GHG emissions in the United States in 2012 totaled

approximately 6,500 MMTCO₂e, of which 5,500 MMTCO₂e came from energy consumption. Our current analysis suggests that ultimate emissions from these leases could account for approximately 21% of total U.S. GHG emissions and 24% of energy-related GHG emissions.

As the tables and figures show, coal extraction was consistently the largest contributor to overall emissions in 2012. Combined, coal and coalbed methane accounted for greater than half of all MTCO₂e. The tables also show the considerable contribution of onshore and offshore natural gas extraction to total emissions of CH₄. This contribution is significant, given the GWP of CH₄; 1 metric ton of CH₄ emitted is equal to approximately 25 metric tons of CO₂ emitted (IPCC, 2007). Table 5 presents coal production and emissions totals by state in 2012.

Indirect methane emissions from natural gas production

Methane emissions from natural gas venting on public lands for which royalties are paid could total almost 1.3 million metric tons CO₂e.

CO₂ emissions from natural gas flaring on public lands for which royalties are paid could total 173,470 metric tons CO₂. These emissions are significantly higher when accounting for emissions from vented and flared natural gas from public lands for which royalties have not been paid.

Source: Appendix to this report.

Table 5. Coal production and emissions by state in 2012

	Type of coal	Sales volume (short ton)	CH ₄ emissions (metric tons)	CO ₂ emissions (metric tons)	N ₂ O emissions (metric tons)	CO ₂ e (metric tons)
Alabama	Bituminous	1,934,725	80	4,634,439	73	4,658,333
Colorado	Bituminous	20,586,124	850	49,311,984	782	49,566,232
Kentucky	Bituminous	207,931	9	498,078	8	500,646
Montana	Sub-bituminous	21,811,626	656	38,701,250	604	38,897,676
New Mexico	Sub-bituminous	4,957,756	149	8,796,747	137	8,841,394
North Dakota	Lignite	3,839,502	86	531,882	79	557,567
Oklahoma	Bituminous	352,171	15	843,590	13	847,940
Utah	Bituminous	13,392,915	553	32,081,377	509	32,246,786
Wyoming	Sub-bituminous	354,972,808	10,683	629,842,604	9,831	633,039,334
Total		422,055,558	13,080	765,241,950	12,037	769,155,909

Note: Totals may not sum due to rounding.

Table 6 presents the total GHG emissions split by onshore and offshore production of fossil fuels. As the table shows, approximately 77% of the emissions evaluated in 2012 came from onshore locations. This was largely driven by the dominance of coal extraction from federal lands, which only occurred onshore. Figure 2 presents the amount of fossil fuels extracted onshore and offshore.

Table 6. GHG emissions (in CO₂e) associated with fossil fuels extracted from federal lands by private leaseholders by location

Fossil fuel	2012 MTCO ₂ e
Onshore	1,028,689,025
Offshore	315,370,364




Figure 2. Percentage of MTCO₂e emissions associated with fossil fuels extracted from federal lands by private leaseholders, by location of extraction.

4. Conclusions

This report presents a preliminary estimate of the magnitude of GHG emissions associated with fossil fuel extraction from federal lands by private leaseholders. Our findings suggest that in 2012, the most recent year for which total U.S. GHG emissions data were available, the estimated 1,344 MMTCO₂e in ultimate downstream GHG emissions from fossil fuel extraction from federal lands by private leaseholders could have accounted for approximately 21% of total U.S. GHG emissions and 24% of energy-related GHG emissions. We include an appendix in this report that presents the results of a quantitative analysis of indirect emissions associated with natural gas and oil production.

References

- Allen, D.T., V.M. Torres, J. Thomas, D.W. Sullivan, M. Harrison, A. Hendler, S.C. Herndon, C.E. Kolb, M.P. Fraser, A.D. Hill, B.K. Lamb, J. Miskimins, R.F. Sawyer, and J.H. Seinfeld. 2013. Measurements of methane emissions at natural gas production sites in the United States. *PNAS USA* 110(44):17768–17773.
- BLM. 2011. 2010 Public Land Statistics: Federal Coal Leases, Table 3-35. U.S. Department of the Interior Bureau of Land Management. Available: http://www.blm.gov/public_land_statistics/pls10/pls3-35_10.pdf. Accessed 6/19/2014.
- BOEMRE. 2014. Outer Continental Shelf Oil and Gas Production. Bureau of Ocean Energy Management, Regulation and Enforcement. Available: http://www.data.bsee.gov/homepg/data_center/production/ocsprod.asp. Accessed 5/1/2014.
- Brandt, AR., G.A. Heath, E.A. Kort, F. O’Sullivan, G. Pétron, S.M. Jordaan, P. Tans, J. Wilcox, A.M. Gopstein, D. Arent, S. Wofsy, N.J. Brown, R. Bradley, G.D. Stucky, D. Eardley, and R. Harriss. 2014. Methane leaks from North American natural gas systems. *Science* 343(6172):733–735.
- Burnham, A., J. Han., C.E. Clark, M. Wang, J.B. Dunn, and I.P. Rivera. 2011. Life-cycle greenhouse gas emissions of shale gas, natural gas, coal and petroleum. *Environ. Sci. Technol.*, doi:10.1021/es201942m.
- Cathles L.M., L. Brown, M. Tamm, and A. Hunter. 2012. A commentary on “The greenhouse-gas footprint of natural gas in shale formations” by R.W. Howarth, R. Santoro, and Anthony Ingraffea. *Climatic Change* 113:525–535. doi: 10.1007/s10584-011-0333-0.

Caulton, D.R., P.B. Shepson, R.L. Santoro, J.P. Sparks, R.W. Howarth, A.R. Ingraffea, M.O.L. Cambaliza, C. Sweeney, A. Karion, K.J. Davis, B.H. Stirm, S.A. Montzka, and B.R. Miller. 2014. Toward a better understanding and quantification of methane emissions from shale gas development. *PNAS USA* 111(17):6237–6242.

CEQ. 2010a. Draft NEPA Guidance on Consideration of the Effects of Climate Change and Greenhouse Gas Emissions. Council on Environmental Quality. February 18. Available: <http://www.whitehouse.gov/sites/default/files/microsites/ceq/20100218-nepa-consideration-effects-ghg-draft-guidance.pdf>. Accessed 6/10/2014.

CEQ. 2010b. Federal Greenhouse Gas Accounting and Reporting Guidance. Council on Environmental Quality. October 6. Available: http://www.whitehouse.gov/sites/default/files/microsites/ceq/ghg_guidance_document_0.pdf. Accessed 6/19/2014.

CEQ. 2011. Greenhouse Gas Emissions Inventory for the Federal Government: 2010 Data. Available: <http://explore.data.gov/Geography-and-Environment/FY2010-Federal-Government-Greenhouse-Gas-Inventory/vzm3-edjq>. Accessed 6/10/2014.

Chandra, V. 2011. Natgas.info. Available: <http://www.natgas.info/html/natgasunitsconversion.html>. Accessed 6/19/2014.

EIA. 2009a. Annual Coal Report – 2009. DOE/EIA-0584(2009). U.S. Energy Information Administration. Available: <http://www.eia.gov/cneaf/coal/page/acr/acr.pdf>. Accessed 6/4/2014.

EIA. 2009b. Annual Energy Review – 2008. U.S. Energy Information Administration. Available: <http://www.eia.gov/FTP/ROOT/multifuel/038408.pdf>. Accessed 6/4/2014.

EIA. 2010. Annual Energy Review – 2009. U.S. Energy Information Administration. Available: <http://205.254.135.24/FTP/ROOT/multifuel/038409.pdf>. Accessed 6/4/2014.

EIA. 2011a. Annual Energy Review – 2010. U.S. Energy Information Administration. Available: <http://205.254.135.24/totalenergy/data/annual/>. Accessed 6/4/2014.

EIA. 2011b. What is Energy? U.S. Energy Information Administration. Available: http://www.eia.gov/kids/energy.cfm?page=about_energy_conversion_calculator-basics. Accessed 6/4/2014.

EIA. 2012a. Annual Energy Outlook 2012 with Projection to 2035. U.S. Energy Information Administration. Available: <http://www.eia.gov/forecasts/archive/aeo12/>. Accessed 6/19/2014.

- EIA. 2012b. Annual Energy Review 2011. U.S. Energy Information Administration. Available: <http://www.eia.gov/totalenergy/data/annual/pdf/aer.pdf>. Accessed 6/19/2014.
- EIA. 2013. Annual Energy Outlook 2013 with Projections to 2040. U.S. Energy Information Administration. Available: <http://www.eia.gov/forecasts/archive/aeo13/>. Accessed 6/19/2014.
- EIA. 2014a. Annual Energy Outlook 2014 with Projections to 2040. U.S. Energy Information Administration. Available: <http://www.eia.gov/forecasts/aeo/>. Accessed 6/19/2014.
- EIA. 2014b. Coal Data: U.S. Coal Production, Annual. U.S. Energy Information Administration. Available: <http://www.eia.gov/coal/data.cfm#production>. Accessed 6/10/2014.
- EIA. 2014c. Natural Gas Data: Natural Gas Gross Withdrawals and Production, Annual. U.S. Energy Information Administration. Available: http://www.eia.gov/dnav/ng/ng_prod_sum_dcu_NUS_a.htm. Accessed 6/11/2014.
- EIA. 2014d. Petroleum and Other Liquids Data: Crude Oil Production, Annual. U.S. Energy Information Administration. Available: http://www.eia.gov/dnav/pet/pet_crd_crpdn_adc_mbbbl_a.htm. Accessed 6/11/2014.
- Hayhoe, K., H.S. Khashgi, A.K. Jain, and D.J. Wuebbles. 2002. Substitution of natural gas for coal: Climatic effects of utility sector emissions. *Climatic Change* 54:107–139.
- Howarth, R.W., R. Santoro, and A. Ingraffea. 2011. Methane and the greenhouse gas footprint of natural gas from shale formations. *Climatic Change Letters* 106:679–690. doi: 10.1007/s10584-011-0061-5.
- Howarth, R.W., R. Santoro, and A. Ingraffea. 2012a. Venting and leakage of methane from shale gas development: Response to Cathles et al. *Climatic Change* 113:537–549. doi: 10.1007/s10584-012-0401-0.
- Howarth, R.W., D. Shindell, R.L. Santoro, A. Ingraffea, N. Phillips, and A. Townsend-Small. 2012b. Methane Emissions, from Natural Gas Systems. National Climate Assessment, February. Report No. 2011-003. Office of Science and Technology Policy, Washington DC.
- Hultman N., D. Rebois, M. Scholten, and C. Ramig. 2011. The greenhouse impact of unconventional gas for electricity generation. *Environ Res Lett* 6:044008, doi:10.1088/1748-9326/6/4/044008.
- IEA. 2011. International Energy Agency Unit Converter. Available: <http://www.i.e.a.org/stats/unit.asp>. Accessed 5/30/2014.

- IPCC. 2006. *IPCC Guidelines for National Greenhouse Gas Inventories, Vol. 2: Energy*. Intergovernmental Panel on Climate Change. Available: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol2.html>. Accessed 5/30/2014.
- IPCC. 2007. Changes in atmospheric constituents and in radiative forcing. Chapter 2 in *IPCC Fourth Assessment Report: Climate Change 2007*. Intergovernmental Panel on Climate Change.
- Jiang M., W.M. Griffin, C. Hendricksin, P. Jaramillo, J. van Briesen, and A. Benkatesh. 2011. Life cycle greenhouse gas emissions of Marcellus shale gas. *Environ Res Lett* 6:03414, doi: 10.1088/1748-9326/6/3/034014.
- Karion, A. 2013. Methane emissions estimate from airborne measurements over a western United States natural gas field. *Journal of Geophys Res* 10.1002/grl.50811.
- Lelieveld, J., S. Lechtenbohmer, S.S. Assonov, C.A.M. Brenninkmeijer, C. Dinest, M. Fishedick, and T. Hanke. 2005. Low methane leakage from natural gas pipelines. *Nature* 434:841–842.
- Miller, S.M., S.C. Wofsy, A.M. Michalak, E.A. Kort, A.E. Andrews, S.C. Biraud, E.J. Dlugokencky, J. Eluszkiewicz, M.L. Fischer, G. Janssens-Maenhout, B.R. Miller, J.B. Miller, S.A. Montzka, T. Nehrkorn, and C. Sweeney. 2013. Anthropogenic emissions of methane in the United States. *PNAS USA* 110(50):20018–20022.
- NETL. 2011. Life Cycle Greenhouse Gas Analysis of Natural Gas Extraction & Delivery in the United States (updated May 23, 2011). National Energy Technology Laboratory. Presented at Cornell University Lecture Series. Available: http://www.netl.doe.gov/energy-analyses/pubs/NG_LC_GHG_PRES_12MAY11.pdf. Accessed 6/3/2014.
- ONRR. 2011. Statistical Information: Reported Royalty Revenues, Sales Volumes, Calendar Years 2008 to 2012. Office of Natural Resources Revenue. Data obtained directly from ONRR in January 2012.
- ONRR. 2014a. Oil and Gas Operations Report. Vented and Flared Gas Volumes (Mcf). Calendar Years 2008–2013. Data obtained directly from the Office of Natural Resources Revenue on August 6, 2014.
- ONRR. 2014b. Statistical Information: Reported Royalty Revenues, Sales Volumes, Calendar Years 2008 to 2012. Office of Natural Resources Revenue. Data obtained directly from ONRR on May 23, 2014. Available: <http://statistics.onrr.gov/>. Accessed 6/3/2014.
- Petron G., G.J. Frost, B.R. Miller, A.I. Hirsch, S.A. Montzka, A. Karion, M.K. Trainer, C. Sweeney, A.E. Andrews, L. Miller, J. Kofler, A. Bar-Ilan, E.J. Dlugokencky, L. Patrick, C.T.

Moore, T.B. Ryerson, C. Siso, W. Kolodzey, P.M. Lang, T.J. Conway, P.C. Novelli, K. Alan Masarie, B.D. Hall, D. Guenther, D.R. Kitzis, J.B. Miller, D.C. Welsh, D.E. Wolfe, W.D. Neff, and P.P. Tans. 2012. Hydrocarbon emissions characterization in the Colorado Front Range – A pilot study. *Journal of Geophysical Research* 10.1029/2011JD016360.

Stephenson T., J.E. Valle, and X. Riera-Palou. 2011. Modeling the relative GHG emissions of conventional and shale gas production. *Environ Sci Tech* 45:10757–10764.

The White House Office of the Press Secretary. 2009. Executive Order. Federal Leadership in Environmental, Energy, and Economic Performance. October 5. Available: http://www.whitehouse.gov/assets/documents/2009fedleader_eo_rel.pdf. Accessed 6/1/2014.

U.S. EPA. 2011. Green Power Equivalency Calculator. U.S. Environmental Protection Agency. Available: <http://www.epa.gov/greenpower/pubs/calculator.htm>. Accessed 6/19/2014.

U.S. EPA. 2014a. 2014 Inventory of Greenhouse Gas Emissions and Sinks. U.S. Environmental Protection Agency. Available: <http://www.epa.gov/climatechange/ghgemissions/usinventoryreport.html>. Accessed 6/11/2014.

U.S. EPA. 2014b. Greenhouse Gas Equivalencies Calculator. U.S. Environmental Protection Agency. Available: www.epa.gov/cleanenergy/energy-resources/calculator.html. Accessed 7/7/2014.

Venkatesh A., P. Jamarillo, W.M. Griffin, and H.S. Matthews. 2011. Uncertainty in life cycle greenhouse gas emissions from United States natural gas end-uses and its effect on policy. *Environ Sci Tech* 45:8182–8189.

Appendix. Indirect GHG Emissions Associated with Natural Gas and Oil Production

Indirect emissions are associated with nearly all stages of fossil fuel extraction, broadly defined as any GHG emissions caused by the exploration, production, refinement, or transportation of fossil fuels that are not otherwise accounted for in the ultimate GHG emissions calculations we describe in Section 2.1.

In this appendix, we quantify indirect GHG emissions¹³ associated with natural gas and oil production including venting, flaring, and fugitive emissions. Our focus is on CH₄ emissions from natural gas and oil production, the second largest contributor to climate change after CO₂. CH₄ is 25 times more potent than CO₂ over a 100-year time horizon (IPCC, 2007).

Significant CH₄ emissions are associated with natural gas and oil production. This is driven by the increase in unconventional gas production methods, such as shale and tight gas, as well as venting and flaring during oil production. But, CH₄ emissions are not well quantified and are worthy of further research. Specifically, characterization and accounting for leaks during natural gas production is subject to ongoing review. For example, EPA has revised CH₄ emissions estimates from the natural gas sector several times in the past few years (U.S. EPA, 2014a).

Venting and fugitive (unintentional) leaks of natural gas emit primarily CH₄, while natural gas flaring emits primarily CO₂. To evaluate indirect GHG emissions associated with natural gas production on federal lands, we present four emissions scenarios:

1. 100% venting, using DOI ONRR data on royalties reported for venting and flaring in natural gas production (ONRR, 2014b)
2. 100% flaring, using ONRR data on royalties reported for venting and flaring in natural gas production (ONRR, 2014b)
3. Venting and flaring for onshore and offshore U.S. natural gas and oil production, using ONRR Oil and Gas Operations Report (OGOR) data (ONRR, 2014a).

13. We have included CH₄ and CO₂ in the indirect emissions estimates. Natural gas also includes nitrogen and non-methane volatile organic compounds, which are not included in the current analysis.

4. CH₄ emissions scenarios from literature review
 - a. Unconventional production, upstream/midstream (well site and processing) CH₄ leakage for low and high scenarios
 - b. Conventional production, upstream/midstream CH₄ leakage for low and high scenarios
 - c. Downstream (pipelines and storage) CH₄ leakage for low and high scenarios.

Natural gas venting and flaring

ONRR collects data on royalties reported for natural gas venting and flaring on public lands. These data do not separate venting and flaring quantities nor do they differentiate between conventional and unconventional natural gas production. Because we do not know the fraction of natural gas that producers vent versus flare, we present two emissions scenarios for the ONRR royalties reported for venting and flaring data: a scenario that assumes 100% venting and a scenario that assumes 100% flaring. We also assume that the venting is occurring in the production, rather than the pipeline, stage.

ONRR also collects reported data on venting and flaring from U.S. natural gas and oil production in its OGOR. These data are separated into venting and flaring categories, but include both federal and some non-federal leases that are part of the same communitization agreement.¹⁴ Because we do not know the fraction of these releases that come directly from federal lands, the estimates provided reflect venting and flaring emissions for both federal and some non-federal sources.

Methodology for calculating venting emissions

The first step in calculating venting emissions was to convert the natural gas (Mcf) into CH₄ and CO₂. For these calculations, we assume that natural gas in the production stage is 78.8% CH₄ and 1.52% CO₂ (NETL, 2011). The second step is converting CH₄ and CO₂ into metric tons. For CH₄, the emissions factor is 0.0193 metric tons CH₄ per Mcf.¹⁵ For CO₂, the emissions factor is 0.0519 metric tons CO₂ per McF.¹⁶ The third step is converting the CH₄ emissions from metric tons into CO₂e, using the GWP factor. One metric ton of CH₄ emitted is equal to approximately 25 metric tons of CO₂ emitted (IPCC, 2007).

14. In most cases a portion of the federal leases are managed as part of communitization agreements, which are a collection of federal and some non-federal leases that draw from the same oil and gas reservoir. A non-federal lease could include an American Indian, state, and/or fee lease.

15. <http://www.epa.gov/cmop/resources/convertter.html#two>.

16. http://www.uigi.com/co2_quantity_convert.html.

We present the CH₄, CO₂, and CO₂e emissions associated with the ONRR royalties reported venting scenario in Table A.1.

Table A.1. CH₄ and CO₂ emissions associated with natural gas venting: ONRR royalties reported data

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	2,571	2,315	1,877	5,924	51,757
CH ₄ -CO ₂ e	64,287	57,879	46,925	148,088	1,293,929
CO ₂	133	120	97	307	2,685

Methodology for calculating flaring emissions

The first step in calculating flaring emissions was to convert natural gas volume into weight. The second step was to apply emissions factors for CO₂ and CH₄ to the natural gas quantities. We used emissions factors from the National Energy Technology Laboratory report: 1 lb of flared natural gas is equivalent to 2.67 lbs CO₂ and 1.53E02 lbs CH₄. We then converted pounds into metric tons. In the third step we converted CH₄ emissions from metric tons into CO₂e, using the GWP factor of 25. We present the CH₄, CO₂, and CO₂e emissions from the ONRR royalties reported flaring scenario in Table A.2.

Table A.2. CH₄ and CO₂ emissions associated with natural gas flaring: ONRR royalties reported data

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	49.39	44.46	36.05	113.77	994.04
CH ₄ -CO ₂ e	1,235	1,112	901	2,844	24,851
CO ₂	8,619	7,760	6,291	19,853	173,470

Table A.3 presents GHG emissions from the venting and flaring data from the OGOR, using the methodology described above. As noted above, the OGOR data include reported volumes of natural gas venting and flaring from both natural gas and oil production, and include both federal and some non-federal leases.

Table A.3. CH₄ and CO₂ emissions from venting and flaring gas in onshore and offshore oil and natural gas production: OGOR data

	2008	2009	2010	2011	2012	2013
CH ₄	74,495	79,860	108,389	149,808	157,200	174,982
CH ₄ -CO ₂ e	1,862,375	1,996,511	2,709,728	3,745,203	3,930,006	4,374,543
CO ₂	2,731,487	2,418,951	2,444,369	2,700,567	3,505,468	3,978,642

CH₄ emissions from literature review: venting and unintentional leaks, or fugitive emissions

The venting and flaring emissions discussed above do not account for unintentional leakage. To provide a more comprehensive characterization of CH₄ emissions that may be associated with natural gas production, including unintentional or “fugitive” leaks, we conducted a review of recent studies. We found a wide range of estimates for CH₄ emissions estimates for all stages of natural gas production. The following tables present CH₄ and CO₂ emissions resulting from scenarios derived through the literature review. We recognize that different ranges could be applied. The CH₄ leakage estimates are applied to ONRR onshore gas production data. We calculate CH₄ and CO₂ emissions for the low and high estimates of CH₄ leakage for upstream/midstream and downstream unconventional (shale and tight gas) and conventional natural gas production.

CH₄ emissions are calculated using either bottom-up or top-down studies. CH₄ emissions for official inventories are estimated from the “bottom up,” in which emissions factors are multiplied by activity data from U.S. GHG inventories (e.g., the EPA GHG Inventory). In contrast, top-down studies use direct CH₄ measurements in natural gas production basins (Petron et al., 2012; Karion, 2013). We include estimates derived from both methodologies in the scenarios below.

Methodology for calculating CH₄ and CO₂ emissions for leakage scenarios

The first step in calculating CH₄ and CO₂ emissions for each of the scenarios is to convert natural gas (Mcf) into CH₄. We assume 78.8% CH₄ for production natural gas (i.e., upstream and midstream) and 93.4% CH₄ for pipeline natural gas (i.e., downstream). We assume that production-quality natural gas is 1.52% CO₂, and pipeline quality natural gas is 0.47% CO₂ (NETL, 2011). The second step is to calculate the fraction of unconventional and conventional natural gas production. This step only applies to upstream and midstream scenarios, not to downstream scenarios.¹⁷ The third step is to convert the resulting CH₄ and CO₂ emissions into metric tons. For CH₄, there are 0.0193 metric tons CH₄ per Mcf. For CO₂, there are 0.0519 metric tons CO₂ per Mcf. The fourth step is to apply the CH₄ leakage rate for a given scenario to the total CH₄ and CO₂ quantities and to calculate the emissions associated with leakage. The final step is to convert the CH₄ emissions into CO₂e, using the GWP factor of 25.

17. For this analysis, we assumed that natural gas produced on federal lands mirrors the U.S. split between conventional and unconventional (shale and tight gas) natural gas production. According to EIA, unconventional gas production has grown from 39% to 61% of total natural gas production between 2008 and 2012 (*Annual Energy Outlook 2013 with Projections to 2040*, EIA, 2013).

We have included bottom-up and top-down estimates of percent CH₄ leakage in natural gas production in Table A.4. As stated above, the range of emissions estimates is significant. Upstream (at the well site) and midstream (at gas processing plants) unconventional CH₄ emissions range from 0.6% to 17.3%.¹⁸

Table A.4. Unconventional gas (% CH₄ produced over lifetime of well)^a

Stephenson et al. (2011)	0.6%
U.S. EPA (2011) ^b	3.0%
Howarth et al. (2011)	3.3%
Caulton et al. (2014)	2.8–17.3%
Petron et al. (2012)	2.3–7.7%
Karion (2013)	6.2–11.7%

a. The CH₄ estimates were reviewed in Howarth et al. (2012b) and Caulton et al. (2014).

b. The EPA estimate in this table is calculated in Howarth et al. (2012a), using national emissions from EPA reports and national gas production data from DOE.

Using the range of emissions estimates from the studies listed above, we calculated the resulting indirect CH₄ and CO₂ emissions from unconventional natural gas produced on public lands for the low and high scenarios. Tables A.5 and A.6 present emissions that may be associated with low and high scenarios for unconventional upstream and midstream natural gas production.

We provide estimates for upstream and midstream emissions from conventional natural gas production below, expressed as the percentage of CH₄ produced over the lifecycle of a well. The estimates in the studies included a range from 0.4% to 2.0%, as we present in Table A.7.

Using the range of emissions estimates from the studies listed above, we calculated the resulting indirect CH₄ and CO₂ emissions that may be associated with conventional natural gas produced on public lands. Table A.8 presents the low scenario and Table A.9 presents the high scenario for upstream and midstream emissions.

18. Other recent studies that are not included in the compiled emissions estimates illustrate the uncertain nature of these estimates. In their 2013 study, Allen et al. (2013) found lower CH₄ emissions than EPA estimates, using scaled actual measurements. The Allen et al. (2013) study covered 190 onshore natural gas sites in the United States. Another study, by Miller et al. (2013), used atmospheric data for the United States; the authors found that EPA inventories underestimated CH₄ emissions. Still another study, Brandt et al. (2014), found that measured natural gas emissions rates were higher than inventory estimates, possibly because of a small number of “superemitting” wells.

Table A.5. CH₄ and CO₂ emissions associated with natural gas production from public lands: Unconventional gas, upstream and midstream, low scenario (0.6% CH₄ leakage)

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	99,408	91,191	108,030	122,172	130,011
CH ₄ -CO ₂ e	2,485,188	2,279,771	2,700,758	3,054,311	3,250,281
CO ₂	5,156	4,730	5,604	6,337	6,744

Table A.6. CH₄ and CO₂ emissions associated with natural gas production from public lands: Unconventional gas, upstream and midstream, high scenario (17.3% CH₄ leakage)

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	2,866,251	2,629,336	3,114,874	3,522,638	3,748,658
CH ₄ -CO ₂ e	71,656,265	65,733,389	77,871,850	88,065,962	93,716,440
CO ₂	148,676	136,387	161,573	182,724	194,448

Table A.7. Conventional gas^a

U.S. EPA (2011)	1.6%
Howarth et al. (2011)	1.4%
Hayhoe et al. (2002)	1.2%
Hultman et al. (2011)	1.3%
Venkatesh et al. (2011)	1.8%
Burnham et al. (2011)	2.0%
Stephenson et al. (2011)	0.4%
Cathles et al. (2012)	0.9%

a. The CH₄ estimates from each study were reviewed in Howarth et al. (2012b).

Table A.8. CH₄ and CO₂ emissions associated with natural gas production from public lands: Conventional gas, upstream and midstream, low scenario (0.4% CH₄ leakage)

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	103,656	77,374	78,022	61,443	55,415
CH ₄ -CO ₂ e	2,591,393	1,934,351	1,950,547	1,536,086	1,385,366
CO ₂	5,377	4,013	4,047	3,187	2,874

Table A.9. CH₄ and CO₂ emissions associated with natural gas production from public lands: Conventional gas, upstream and midstream, high scenario (2.0% CH₄ leakage)

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	518,279	386,870	390,109	307,217	277,073
CH ₄ -CO ₂ e	12,956,965	9,671,755	9,752,736	7,680,431	6,926,829
CO ₂	26,884	20,067	20,235	15,936	14,372

Recent estimates in peer-reviewed literature for downstream emissions of CH₄ from natural gas systems range from 0.07% to 10% of the CH₄ produced over the lifetime of the well.

Downstream emissions refer to emissions that occur in storage, and transmission and distribution pipelines. Table A.10 presents the range of downstream CH₄ emissions from various studies.

Table A.10. Downstream CH₄ emissions^a

Hayhoe et al. (2002)	0.2–10%
Lelieveld et al. (2005)	1.0–2.5%
Howarth et al. (2011)	1.4–3.6%
U.S. EPA (2011)	0.9%
Jiang et al. (2011)	0.4%
Hultman et al. (2011)	0.9%
Venkatesh et al. (2011)	0.4%
Burnham et al. (2011)	0.6%
Stephenson et al. (2011)	0.07%
Cathles et al. (2012)	0.7%

a. The CH₄ estimates from each study were reviewed in Howarth et al. (2012b) and Caulton et al. (2014).

Tables A.11 and A.12 present the low and high scenarios for CH₄ and CO₂ emissions for downstream emissions, respectively.

Table A.11. Associated CH₄ and CO₂ emissions from public lands: Downstream^a emissions from natural gas, low scenario (0.07% CH₄ leakage)

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	35,247	28,659	31,122	29,639	29,473
CH ₄ -CO ₂ e	881,175	716,483	778,058	740,979	736,815
CO ₂	477	388	421	401	399

a. Downstream emissions occur in transmission and distribution pipelines and in storage.

Table A.12. Associated CH₄ and CO₂ emissions from public lands: Downstream emissions from natural gas, high scenario (10% CH₄ leakage)

Emissions (metric tons)	2008	2009	2010	2011	2012
CH ₄	5,035,286	4,094,189	4,446,004	4,234,165	4,210,372
CH ₄ -CO ₂ e	125,882,140	102,354,727	111,151,111	105,854,124	105,259,307
CO ₂	68,137	55,402	60,164	57,297	56,975

